



# Minimum number of affine simplices of given dimension



István Szalkai<sup>a,\*</sup>, Zsolt Tuza<sup>b,c</sup>

<sup>a</sup> Department of Mathematics, University of Pannonia, Veszprém, Hungary

<sup>b</sup> Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences, Budapest, Hungary

<sup>c</sup> Department of Computer Science and Systems Technology, University of Pannonia, Veszprém, Hungary

## ARTICLE INFO

### Article history:

Received 26 September 2013

Received in revised form 19 July 2014

Accepted 22 July 2014

Available online 12 August 2014

### Keywords:

Linear hypergraph

Extremal set theory

Euclidean affine simplex

Minimal linear dependency

Stoichiometry

## ABSTRACT

In this paper we formulate and solve extremal problems in the Euclidean space  $\mathbb{R}^d$  and further in hypergraphs, originating from problems in stoichiometry and elementary linear algebra. The notion of affine simplex is the bridge between the original problems and the presented extremal theorem on set systems. As a sample corollary, it follows that if no triple is collinear in a set  $S$  of  $n$  points in  $\mathbb{R}^3$ , then  $S$  contains at least  $\binom{n}{4} - cn^3$  affine simplices for some constant  $c$ . A function related to Sperner's Theorem and its well-known extension to reciprocal sums is also considered and its relation to Turán's hypergraph problems is discussed.

© 2014 Elsevier B.V. All rights reserved.

## 1. Introduction

The roots of the present study date back to the mid-1980s, to the paper by S. Kumar and Á. Pethő [7], concerning an application of linear algebra in stoichiometry. From the algebraic point of view, their very natural question asks about the number of those subsets of a set of vectors which are linearly dependent but each of whose proper subsets is independent. Here we give an asymptotically tight solution for the minimum in terms of dimension and the number of vectors when low-dimensional dependencies are excluded. Our method is to prove a more general result in extremal set theory (Theorem 6 below), hence without assuming anything about the structure of algebraic dependencies.

### 1.1. Motivation in chemistry

Restricting attention to a “universe” of  $D$  kinds of atoms (or atomic parts), each molecule (species) can be represented with a vector in  $\mathbb{R}^D$  whose  $i$ th coordinate means the number of atoms of  $i$ th type in the molecule in question.<sup>1</sup> Then a chemical reaction naturally corresponds to a zero-sum linear combination of these vectors (using the law of mass balance).

The reaction is called minimal if none of the molecules, taking role in it, can be omitted so that the remaining ones could form still a (nother) reaction. In the language of linear algebra this assumption is equivalent to the property that the corresponding set of vectors is linearly dependent but each proper subset of it is independent, that is the defining condition of *linear algebraic simplex*. Both from practical and theoretical purposes the following problem was raised:

\* Corresponding author. Tel.: +36 88624227; fax: +36 88624227.

E-mail addresses: [szalkai@almos.uni-pannon.hu](mailto:szalkai@almos.uni-pannon.hu) (I. Szalkai), [tuza@dcs.uni-pannon.hu](mailto:tuza@dcs.uni-pannon.hu) (Z. Tuza).

<sup>1</sup> The types of atoms are supposed to be in a fixed order. E.g., if  $D = 3$  and the universe is [C, H, O], then we have the vector (0, 2, 1) for H<sub>2</sub>O and (2, 4, 2) for CH<sub>3</sub>COOH.

**Problem 1.** What is the minimum and maximum number of linear algebraic simplices  $S \subseteq \mathcal{V}$  in a set  $\mathcal{V}$  of vectors in  $\mathbb{R}^D$  if only the size  $|\mathcal{V}|$  is given and  $\mathcal{V}$  spans  $\mathbb{R}^D$ ? What are the structures of sets  $\mathcal{V}$  which contain extremal number of simplices?

The answer was given in [8]. Moreover, Problem 1 was generalized for matroids in [4]; actually its authors solved it a decade earlier than published, see [3].

Concerning the minimum, the results in [8] show that almost all vectors must be *parallel*, i.e. almost all molecules (species) are isomer molecules or multiple doses. The problem where parallel vectors are excluded is still unsolved in general:

**Problem 2.** What is the minimum number of linear algebraic simplices  $S \subseteq \mathcal{V}$  if only the size  $|\mathcal{V}|$  is given,  $\mathcal{V}$  does not contain parallel vectors and  $\mathcal{V}$  spans  $\mathbb{R}^D$ ? What are the structures of sets  $\mathcal{V}$  which contain the minimum number of simplices?

A conjecture on both the minimum number and the structure attaining it is stated in [9]. The cases  $D = 3$  and  $D = 4$  were solved in [9,17], respectively.

### 1.2. Geometric formulation

In the framework of linear algebra the problem is somewhat non-symmetric because the zero vector plays a special role. This asymmetry can be eliminated if we translate the problem to the language of geometry. Moreover, restricting attention to sets  $\mathcal{V} \subset \mathbb{R}^D$  containing neither the zero vector nor a pair of parallel vectors, the dimension can be reduced from  $D$  to  $d = D - 1$ : first associate each element  $v \in \mathcal{V}$  with its direction  $\lambda \cdot v$  ( $\lambda \in \mathbb{R}$ ), and then intersect this system  $\lambda \mathcal{V}$  with a  $(D - 1)$ -dimensional hyperplane  $\mathcal{P}$  which does not contain the origin and is not parallel to any element of  $\mathcal{V}$ .

The mapping from  $\mathcal{V}$  to the set  $\mathcal{V}^{\mathcal{P}} := \lambda \mathcal{V} \cap \mathcal{P}$  is a bijection under which linear algebraic simplices  $S \subset \mathcal{V}$  correspond to *affine simplices*  $S^{\mathcal{P}} \subset \mathbb{R}^{D-1}$ , where a set  $S$  of  $k \geq 3$  points in the Euclidean  $d$ -space is called an *affine simplex* if  $S$  is contained in some  $(k - 2)$ -dimensional hyperplane but no proper subset  $S' \subsetneq S$  is contained in a hyperplane of dimension  $|S'| - 2$ . For instance, in  $\mathbb{R}^3$  the following three types of affine simplices occur:

- three collinear points;
- four coplanar points, no three of which are collinear;
- five points, no four of which are coplanar.

Affine simplices can alternatively be defined by requiring that the vectors  $s_2 - s_1, s_3 - s_1, \dots, s_k - s_1$  be linearly dependent but their proper subsets should not (for every choice of a point to be labeled  $s_1$ ).

In cases of low dimension, as solved in [9,17], almost all points of the extremal configurations for Problem 2 attaining the minimum number of affine simplices lie on one or two lines, i.e. mostly contain affine simplices of three points. In this way the natural question arises to determine the minimum in the other extreme, where no three points are collinear. For this reason our goal is to study point sets which contain no affine simplices smaller than a given size. The first interesting case is  $\mathbb{R}^3$ .

Let  $S \subset \mathbb{R}^d$  be a set of  $n$  points, no  $d$  of which lie on a  $(d - 2)$ -dimensional hyperplane. Then two kinds of subsets of  $S$  form an affine simplex:

- $d + 1$  points on a hyperplane of dimension  $d - 1$ , or
- $d + 2$  points, no  $d + 1$  of which lie on a common hyperplane of dimension  $d - 1$ .

**Theorem 3.** For every  $d \geq 3$  there is a constant  $c = c(d)$  with the following property. If  $S \subset \mathbb{R}^d$  is a set of  $n$  points, no  $d$  of them lying on a hyperplane of dimension  $d - 2$ , then  $S$  determines at least  $\binom{n}{d+1} - cn^d$  affine simplices.

**Corollary 4.** For any  $n$  points in the 3-space, no three being collinear, the number of coplanar quadruples plus the 5-tuples containing no coplanar quadruples is at least  $\binom{n}{4} - O(n^3)$  as  $n \rightarrow \infty$ .

These results are asymptotically tight, as shown by the obvious example of  $n$  coplanar points in  $\mathbb{R}^3$  (no three of them being on a line) and also for any  $d \geq 3$  by  $n$  points of  $\mathbb{R}^{d-1}$  in general position when embedded isometrically into  $\mathbb{R}^d$ . Such a set of points has exactly  $\binom{n}{d+1}$  affine simplices. In fact, configurations with even fewer affine simplices exist, which in addition span the  $d$ -space. For instance,  $n - 1$  points of  $\mathbb{R}^{d-1}$  in general position embedded in a hyperplane of  $\mathbb{R}^d$  plus an  $n$ th point outside that hyperplane generate just  $\binom{n-1}{d+1}$  affine simplices (as no affine simplex contains the  $n$ th point).

In  $\mathbb{R}^3$ , the two arrangements of points just mentioned yield  $\frac{1}{24}n^4 - \frac{1}{4}n^3 + O(n^2)$  and  $\frac{1}{24}n^4 - \frac{5}{12}n^3 + O(n^2)$ , respectively. Currently we do not know whether or not the latter error term  $\frac{5}{12}n^3$  is asymptotically tight. We do know, however, that the construction above is not extremal; an improvement of the order  $O(n^2)$  will be proved in Proposition 7.

### 1.3. Combinatorial formulation

Here we put the problems and results above in a more general setting. Let  $\mathcal{H} = (X, \mathcal{E})$  be a hypergraph, where  $X$  is the finite vertex set and  $\mathcal{E}$  is the edge set consisting of subsets of  $X$ . We extend the notion of linear hypergraph (also called “simple” or “almost disjoint” in some parts of the literature) as follows.

**Definition 5.** We say that a hypergraph  $\mathcal{H} = (X, \mathcal{E})$  is  $q$ -linear (for some integer  $q \geq 1$ ) if  $|E \cap E'| < q$  holds for all  $E, E' \in \mathcal{E}, E \neq E'$ .

Hence, in a 1-linear hypergraph any two edges are disjoint, and 2-linear coincides with linear hypergraphs in the usual sense, in analogy with Euclidean spaces where any two points uniquely determine a line.

We also introduce some notation. As usual,  $\binom{S}{k}$  will stand for the collection of all  $k$ -element subsets of set  $S$ . For any hypergraph  $\mathcal{H} = (X, \mathcal{E})$ , let

- $\mathcal{E}_k := \bigcup_{E \in \mathcal{E}} \binom{E}{k}$ —usually  $(X, \mathcal{E}_k)$  is called the  $k$ -section hypergraph of  $\mathcal{H}$ ;
- $\mathcal{E}_{k+1}^0 := \{F \in \binom{X}{k+1} \mid \binom{F}{k} \cap \mathcal{E}_k = \emptyset\}$ .

Corresponding to  $k = d + 1$ , in analogy with the geometric interpretation, we call the members of  $\mathcal{E}_k \cup \mathcal{E}_{k+1}^0$  the  $(k - 1)$ -dimensional *semi-simplices*<sup>2</sup> in  $\mathcal{H}$ .

**Theorem 6.** For every  $k \geq 3$  there is a constant  $c = c(k)$  such that

$$|\mathcal{E}_k| + |\mathcal{E}_{k+1}^0| \geq \binom{n}{k} - cn^{k-1}$$

holds for all  $(k - 1)$ -linear hypergraphs  $\mathcal{H} = (X, \mathcal{E})$  on  $n$  vertices.

This result implies [Theorem 3](#), by considering the hypergraph whose edges are the sets of points lying on a common hyperplane of dimension  $d - 1$ .

#### 1.4. Sperner families and Turán numbers

For any hypergraph  $\mathcal{H} = (X, \mathcal{E})$  (not necessarily  $q$ -linear for a prescribed value of  $q$ ) and for any  $k$ , the set system  $\mathcal{S}_k(\mathcal{H}) := \mathcal{E}_k \cup \mathcal{E}_{k+1}^0$  is a *Sperner family*, which means that none of its members contains any other.

The well known YBLM inequality<sup>3</sup> states that

$$\sum_{S \in \mathcal{S}} \binom{n}{|S|}^{-1} \leq 1 \tag{1}$$

holds for every Sperner family  $\mathcal{S}$  (where  $n$  is the number of vertices). In particular, (1) is valid for the family  $\mathcal{S} = \mathcal{S}_k(\mathcal{H})$  of any  $\mathcal{H}$ , too. In connection with the main problem studied here, one may also consider the values

$$s(n, k) := \min_{\mathcal{H} \text{ is } (k-1)\text{-linear, } |X|=n} \sum_{S \in \mathcal{S}_k(\mathcal{H})} \binom{n}{|S|}^{-1}$$

and analogously, without assuming  $(k - 1)$ -linearity,

$$s'(n, k) := \min_{\mathcal{H}=(X, \mathcal{E}), |X|=n} \sum_{S \in \mathcal{S}_k(\mathcal{H})} \binom{n}{|S|}^{-1}.$$

Since there exist only finitely many hypergraphs on any given number  $n$  of vertices, both  $s(n, k)$  and  $s'(n, k)$  are well-defined and are at most 1 by the YBLM inequality, for all  $n$  and  $k$ . In [Theorem 8](#) we prove that for every fixed  $k$ , the values of  $s(n, k)$  and  $s'(n, k)$  tend to constants larger than 0 and smaller than 1 as  $n$  gets large. We also consider their relation to the Turán problem on graphs and uniform hypergraphs.

## 2. Proof of the general lower bound

Here we prove [Theorem 6](#). By the free choice of  $c = c(k)$ , we may restrict ourselves to  $n$  sufficiently large, say  $n > k^3$ . Moreover, due to the nature of the problem, we may also assume without loss of generality that  $|E| \geq k$  holds for all  $E \in \mathcal{E}$ . Let  $H \in \binom{X}{k}$  be any  $k$ -tuple. If it is contained in some  $E \in \mathcal{E}$ , then  $H$  is counted in  $\mathcal{E}_k$  precisely once. We will prove that, with

<sup>2</sup> The definition of affine simplices requires that no subset of a simplex can be another simplex (see e.g. [17, Definition 10]), whereas for the members of  $\mathcal{E}_k \cup \mathcal{E}_{k+1}^0$  we require non-containment only for subsets smaller by one element.

<sup>3</sup> For several decades, it was called LYM inequality, stated and proved in exactly that form independently by Yamamoto [20], Meshalkin [11] and Lubell [10] (in this order of chronology). Bollobás [1] proved a more general result, however, from which the inequality follows immediately. Inequalities of this kind have lots of applications in extremal problems in various areas of mathematics; cf. the two-part survey [18,19]. The current acronym YBLM coincides (apart from punctuation) with the abbreviated name of famous Hungarian architect Miklós Ybl (1814–1891) [21].

possibly few exceptions, also the other  $k$ -tuples  $H$  generate at least one member of  $\mathcal{E}_{k+1}^0$  on the average. More explicitly, it will turn out that most of those sets  $H$  can be completed to a member of  $\mathcal{E}_{k+1}^0$  in more than  $k$  different ways.

From now on we assume that  $H \in \binom{X}{k} \setminus \mathcal{E}_k$ . Let  $x \in H$  be any vertex. If the subset  $H \setminus \{x\}$  is contained in an edge of  $\mathcal{H}$ , we denote that edge by  $E_x$ ; and otherwise we define  $E_x := H \setminus \{x\}$ . A more precise and unambiguous notation would be  $E_x(H)$ , but for simplicity we write  $E_x$  as long as just one  $H$  is considered. Note that  $E_x$  is unique for each  $x \in H$  (once  $H$  is understood), since  $\mathcal{H}$  is  $(k - 1)$ -linear. It also follows for any two distinct  $x, x' \in H$  that  $E_x$  and  $E_{x'}$  share no vertex outside  $H$ . We set

$$H^* := \bigcup_{x \in H} E_x.$$

Then we have the implication

$$H \in \binom{X}{k} \setminus \mathcal{E}_k \wedge z \in X \setminus H^* \implies H \cup \{z\} \in \mathcal{E}_{k+1}^0 \tag{2}$$

because  $|(H \cup \{z\}) \cap E'| = k$  for an  $E' \in \mathcal{E}$  with  $z \in E'$  would imply  $E_x = E'$  for some  $x \in H$ , thus  $z \in H^*$  would hold.

We say that the  $k$ -tuple  $H$  is a *near-cover* of  $\mathcal{H}$  if  $|H^*| \geq |X| - k$ . The proof now splits into two situations, whether  $\mathcal{H}$  has, or does not have, a near-cover.

Suppose first that no  $H \in \binom{X}{k} \setminus \mathcal{E}_k$  is a near-cover of  $\mathcal{H}$ . Then by (2) we obtain that each  $H$  can be extended to a member  $F$  of  $\mathcal{E}_{k+1}^0$  in at least  $k + 1$  different ways. On the other hand, each  $F \in \mathcal{E}_{k+1}^0$  can be obtained from exactly  $k + 1$  sets  $H \in \binom{X}{k} \setminus \mathcal{E}_k$ , namely from its  $k$ -element subsets. Thus, in this case we have

$$|\mathcal{E}_{k+1}^0| \geq \left| \binom{X}{k} \setminus \mathcal{E}_k \right|$$

and the inequality stated in the theorem holds even without the error term  $O(n^{k-1})$ .

Suppose now that some  $H \in \binom{X}{k} \setminus \mathcal{E}_k$  is a near-cover of  $\mathcal{H}$ . Then the cardinality of the set

$$X'^* \setminus H$$

is at least  $n - 2k$ , and  $X'$  is partitioned into sets of type  $E'_x := E_x \setminus H$  ( $x \in H$ ). Say,  $X' = E'_{x_1} \cup \dots \cup E'_{x_\ell}$  where  $\ell \leq k$ .

A case that can directly be settled is when some  $\ell - 1$  sets from  $\{E'_{x_1}, \dots, E'_{x_\ell}\}$  cover together at most  $k^3 - 4k^2 + 5k$  vertices. There are at most  $2k$  vertices outside  $X'$ , hence some  $E'_{x_i}$  contains at least  $n - k^3 + 4k^2 - 7k$  elements. Then we obtain that  $|\mathcal{E}_k| \geq \binom{n - k^3 + 4k^2 - 7k}{k} = \binom{n}{k} - O(n^{k-1})$  is valid.<sup>4</sup> Therefore, we may suppose for the rest of the proof that the union of any  $\ell - 1$  sets from  $\{E'_{x_1}, \dots, E'_{x_\ell}\}$  has cardinality greater than  $k^3 - 4k^2 + 5k$ .

Consider any  $H' \in \binom{X'}{k} \setminus (\mathcal{E}_k \cup \{H\})$ . A coincidence  $E_{y_0}(H') = E'_{x_i}$  can happen with only one vertex  $y_0 \in H$  and only one index  $i$  ( $1 \leq i \leq \ell$ ), namely when  $|H' \cap E'_{x_i}| = k - 1$ . If this situation occurs, assume that  $E_{y_0}(H') = E'_{x_1}$ . Then, since  $\mathcal{H}$  is  $(k - 1)$ -linear, for any  $y \in H' \setminus \{y_0\}$  and for any  $i' \neq i$  with  $i' > 1$  we have  $|E_y(H') \cap E'_{x_{i'}}| \leq k - 2$ , and so  $E_y(H')$  meets  $E'_{x_2} \cup \dots \cup E'_{x_\ell}$  in at most  $(\ell - 1)(k - 2) \leq (k - 1)(k - 2)$  vertices, one of which is  $y_0$ . Therefore the  $k - 1$  choices of  $y \neq y_0$  cover at most  $(k^3 - 3k^2 + k) - (k^2 - 3k + 1) + 1 = k^3 - 4k^2 + 4k$  vertices of  $E'_{x_2} \cup \dots \cup E'_{x_\ell}$ . Hence, the inequality

$$\left| \left( E'_{x_2} \cup \dots \cup E'_{x_\ell} \right) \setminus \bigcup_{y \in H'} E_y(H') \right| \geq k + 1$$

follows by  $|X'| \geq n - 2k$  and by the assumed lower bound on  $|E'_{x_2} \cup \dots \cup E'_{x_\ell}|$ . Thus, in this case,  $H'$  can be completed to a member of  $\mathcal{E}_{k+1}^0$  in at least  $k + 1$  different ways. The situation is even better if we have  $|E_y(H') \cap E'_{x_i}| \leq k - 2$  for all  $1 \leq i \leq \ell$ . Then summing over all  $y \in H'$  and all  $1 \leq i \leq \ell$ , we obtain the upper bound

$$|(H'^* \setminus H^* \setminus H)| \leq k^2(k - 2),$$

so that there are at least  $|X' \setminus H'^*| \geq n - (k^3 - 2k^2 + 3k) \geq k + 1$  ways to extend  $H'$  to a member of  $\mathcal{E}_{k+1}^0$  whenever  $n > k^3$  (and  $k \geq 2$ ). Consequently,

$$|\mathcal{E}_{k+1}^0| \geq \left| \binom{X'}{k} \setminus \mathcal{E}_k \right|$$

holds, and therefore  $|\mathcal{E}_k| + |\mathcal{E}_{k+1}^0| \geq \binom{n}{k} - O(n^{k-1})$  is valid as  $n$  gets large, because  $|X'| \geq n - 2k$ .  $\square$

<sup>4</sup> We may actually write the somewhat larger value  $\binom{n - k^3 + 4k^2 - 7k + k - 1}{k}$ , by considering  $E_{x_i}$  instead of  $E'_{x_i}$ ; but this is irrelevant concerning the current proof.

### 3. Geometric upper bound

As we mentioned in the introduction,  $n$  points in  $\mathbb{R}^d$  may generate as few as  $\binom{n-1}{d+1}$  affine simplices, each of which has more than  $d$  points. Focusing on the case  $r = 3$ , here we show that the number of affine simplices can be even smaller in  $\mathbb{R}^3$ .

**Proposition 7.** *There is an arrangement of  $n$  points in  $\mathbb{R}^3$  which determines the following number of affine simplices:*

- $\binom{n-1}{4} - \frac{(n-2)(n-5)}{2}$  if  $n$  is even,
- $\binom{n-1}{4} - \frac{(n-3)(n-5)}{2}$  if  $n$  is odd;

that is,  $\frac{1}{24}n^4 - \frac{5}{12}n^3 + O(n^2)$ .

**Proof.** First, let  $n$  be even. Take  $n - 2$  points  $x_1, \dots, x_{n-2}$  on a plane  $P \subset \mathbb{R}^3$ , such that no three of them are collinear, moreover all the  $n/2 - 1$  lines  $\overline{x_{2i-1}x_{2i}}$  are parallel for  $i = 1, 2, \dots, n/2 - 1$ . Let  $x_{n-1}$  and  $x_n$  be two points outside  $P$ , such that the line  $\overline{x_{n-1}x_n}$  is parallel to  $\overline{x_1x_2}$  (and hence to the other pairs as well). We have the following types of affine simplices:

- quadruples of points in  $P$ ;
- quadruples of the form  $\{x_{2i-1}, x_{2i}, x_{n-1}, x_n\}$  ( $i = 1, 2, \dots, n/2 - 1$ );
- quintuples of the form  $\{x_a, x_b, x_c, x_{n-1}, x_n\}$  ( $1 \leq a < b < c \leq n - 2$ ), where  $\{2i - 1, 2i\} \not\subset \{a, b, c\}$  for any  $i$ .

The number of sets of those three types is  $\binom{n-2}{4}$ ,  $\frac{1}{2}(n - 2)$ , and  $\frac{1}{6}(n - 2)(n - 4)(n - 6)$ , respectively.

If  $n$  is odd, we take  $(n - 3)/2$  pairs of points inside  $P$  which determine lines  $\overline{x_{2i-1}x_{2i}}$  parallel to  $\overline{x_{n-1}x_n}$ , plus one point  $x_{n-2}$  of  $P$  which is not collinear with any two of  $x_1, \dots, x_{n-3}$ . Then we have  $\binom{n-2}{4}$  affine simplices inside  $P$ , further  $\frac{1}{2}(n - 3)$  ones of the form  $\{x_{2i-1}, x_{2i}, x_{n-1}, x_n\}$ , moreover  $\frac{1}{6}(n - 3)(n - 5)(n - 7)$  of the form  $\{x_a, x_b, x_c, x_{n-1}, x_n\}$  ( $1 \leq a < b < c \leq n - 3$ ) where  $\{2i - 1, 2i\} \not\subset \{a, b, c\}$ , and finally  $\frac{1}{2}(n - 3)(n - 5)$  of the form  $\{x_a, x_b, x_{n-2}, x_{n-1}, x_n\}$  not containing any pair  $\{x_{2i-1}, x_{2i}\}$ .  $\square$

### 4. The YBLM inequality

Recall from Section 1.4 that  $s(n, k)$  and  $s'(n, k)$  are defined as the minimum of the sum  $\sum_{S \in \mathcal{S}_k(\mathcal{H})} \binom{n}{|S|}^{-1}$  where  $\mathcal{H}$  runs over all hypergraphs of order  $n$  – with or without assuming  $(k - 1)$ -linearity – and  $\mathcal{S}_k(\mathcal{H}) = \mathcal{E}_k \cup \mathcal{E}_{k+1}^0$ . Here we study the asymptotic behavior of these two functions, and point out a relation to Turán numbers.

#### 4.1. The limits of $s(n, k)$ and $s'(n, k)$

Our goal in this subsection is to prove the following result.

**Theorem 8.** *For every fixed  $k \geq 2$ , the limits*

$$s_k := \lim_{n \rightarrow \infty} s(n, k) \quad \text{and} \quad s'_k := \lim_{n \rightarrow \infty} s'(n, k)$$

exist and satisfy

$$0 < s'_k \leq s_k < 1$$

with strict inequality at both ends. More explicitly,  $\frac{1}{k+1} \leq s'_k \leq s_k \leq 1 - \frac{k}{2^k}$ .

We state three assertions below which together will immediately imply the validity of the theorem as the middle inequality holds by definition.

**Lemma 9.** *For every fixed  $k$ , the sequences  $(s(n, k))_{n=k+1}^\infty$  and  $(s'(n, k))_{n=k+1}^\infty$  are non-decreasing.*

**Proof.** For any hypergraph  $\mathcal{H} = (X, \mathcal{E})$  on  $n$  vertices, let us introduce the notation  $m_k := |\mathcal{E}_k|$  and  $m_{k+1} := |\mathcal{E}_{k+1}^0|$ . The inequality

$$\frac{m_k}{\binom{n}{k}} + \frac{m_{k+1}}{\binom{n}{k+1}} \geq b \tag{3}$$

is equivalent to

$$m_{k+1} \geq b \cdot \binom{n}{k+1} - \frac{n-k}{k+1} m_k \tag{4}$$

for any  $b > 0$ . We are going to prove that if the analogue of (4) is valid for every hypergraph on  $n - 1$  vertices, then it is valid for  $\mathcal{H}$  as well. In what follows, assume that it is valid for  $n - 1$ .

Let  $x \in X$  be any vertex. We derive the hypergraph  $\mathcal{H}_{-x}$  from  $\mathcal{H}$  by removing  $x$  from all edges and deleting the edges which become smaller than  $k$  after cutting out  $x$ . Note that  $\mathcal{H}_{-x}$  is  $q$ -linear whenever  $\mathcal{H}$  is (no matter what  $q$  we choose), although the inverse implication is not valid in general.

Let us denote by  $m_k^x$  and  $m_{k+1}^x$  the values corresponding to  $m_k$  and  $m_{k+1}$  in  $\mathcal{H}_{-x}$ , and by  $\mathcal{E}_{k[-x]}$  and  $\mathcal{E}_{k+1[-x]}^0$  the corresponding families of sets, respectively. We then have

$$m_{k+1}^x \geq b \cdot \binom{n-1}{k+1} - \frac{n-k-1}{k+1} m_k^x \tag{4_x}$$

by assumption.

A  $k$ -element set  $F$  occurs in  $\mathcal{E}_{k[-x]}$  if and only if it is contained in some edge of  $\mathcal{H}_{-x}$ ; and this happens precisely when  $x \notin F$  and some  $E \in \mathcal{E}$  contains  $F$  as a subset. Thus, each  $F \in \mathcal{E}_k$  gives rise to a member of  $\mathcal{E}_{k[-x]}$  for exactly  $n - k$  choices of  $x$ , and no more sets occur in  $\bigcup_{x \in X} \mathcal{E}_{k[-x]}$ . Similarly, each  $F \in \mathcal{E}_{k+1}^0$  yields a member of  $\mathcal{E}_{k+1[-x]}^0$  for exactly  $n - k - 1$  choices of  $x$ , and these are all the sets in  $\bigcup_{x \in X} \mathcal{E}_{k+1[-x]}^0$ . As a consequence, the equalities

$$\sum_{x \in X} m_k^x = (n - k) \cdot m_k \quad \text{and} \quad \sum_{x \in X} m_{k+1}^x = (n - k - 1) \cdot m_{k+1} \tag{5}$$

hold. Thus, summing up (4<sub>x</sub>) for all  $x \in X$  we obtain

$$(n - k - 1) \cdot m_{k+1} \geq bn \cdot \binom{n-1}{k+1} - \frac{n-k-1}{k+1} (n - k) \cdot m_k$$

which is equivalent to (4). This completes the proof.

As one of our referees remarked, an alternative way of performing the inductive calculation is to view  $m_k / \binom{n}{k}$  as the probability that a randomly chosen  $k$ -tuple is contained in some edge of  $\mathcal{H}$ . It is left to the reader to verify that the equations in (5) imply (3) from  $n - 1$  to  $n$ . We thank the referee for the useful conceptual comment.  $\square$

**Lemma 10.** For every fixed  $k$ , we have  $s(k + 1, k) = s'(k + 1, k) = \frac{1}{k+1}$ .

**Proof.** The minimum for both  $s(k + 1, k)$  and  $s'(k + 1, k)$  is attained by the hypergraph with  $k + 1$  vertices and with precisely one edge of cardinality  $k$ .  $\square$

**Lemma 11.** For every fixed  $k$ , we have  $s_k \leq 1 - \frac{k}{2k}$ .

**Proof.** To simplify notation, we consider  $2n$  vertices instead of  $n$ . Consider the hypergraph whose edge set  $\mathcal{E}$  consists of just two disjoint sets of cardinality  $n$  each. It is linear, of course. Moreover, we clearly have

$$|\mathcal{E}_k| = 2 \binom{n}{k} \quad \text{and} \quad |\mathcal{E}_{k+1}^0| = \binom{2n}{k+1} - 2 \binom{n}{k+1} - 2n \binom{n}{k}$$

because a  $(k + 1)$ -tuple does not belong to  $\mathcal{E}_{k+1}^0$  if and only if either it is contained in one of the two edges or it meets one of the edges in precisely one vertex and the other edge in  $k$  vertices. Thus,

$$\begin{aligned} s(2n, k) &\leq \frac{2 \binom{n}{k}}{\binom{2n}{k}} + 1 - \frac{1}{\binom{2n}{k+1}} \left( 2 \binom{n}{k+1} + 2n \binom{n}{k} \right) \\ &= 1 - \frac{2}{\binom{2n}{k}} \left( \frac{k+1}{2n-k} \left( n \binom{n}{k} + \binom{n}{k+1} \right) - \binom{n}{k} \right) \\ &= 1 - \frac{\binom{n}{k}}{\binom{2n}{k}} \cdot 2 \cdot \left( \frac{n(k+1)}{2n-k} + \frac{n-k}{2n-k} - 1 \right) \\ &= 1 - \frac{\binom{n}{k}}{\binom{2n}{k}} \cdot \frac{2nk}{2n-k}. \end{aligned}$$

The function in the last line clearly tends to  $1 - \frac{k}{2k}$  as  $n \rightarrow \infty$ , therefore  $s_k$  cannot be larger.  $\square$

#### 4.2. Turán numbers

For a fixed  $k$ -uniform hypergraph  $\mathcal{F}$ , we use the standard notation  $\text{ex}(n, \mathcal{F})$  for its Turán number; that means the maximum number of edges in a  $k$ -uniform hypergraph of order  $n$  which does not contain any subhypergraph isomorphic to  $\mathcal{F}$ . Further, let  $\mathcal{K}_{k+1}^{(k)}$  denote the hypergraph with  $k + 1$  vertices and  $k + 1$  edges of  $k$  vertices each (i.e., the complete  $k$ -uniform

hypergraph of order  $k + 1$ ). If  $k = 2$  then  $\mathcal{K}_3^{(2)}$  is just the triangle  $K_3$ , the complete graph of order 3. In this very simple case the equality  $\text{ex}(n, K_3) = \lfloor n^2/4 \rfloor$  is well known to hold, but for larger  $k$  the determination of  $\text{ex}(n, \mathcal{K}_{k+1}^{(k)})$  is a famous open problem in extremal hypergraph theory. Extensive surveys on Turán-type problems are, e.g., [6,15]. In particular, a recent summary of results on  $\text{ex}(n, \mathcal{K}_{k+1}^{(k)})$  can be found in the second part of Section 9 in [6].

**Remark 12.** If  $\mathcal{H} = (X, \mathcal{E})$  is a  $k$ -uniform hypergraph of order  $n$  such that each  $(k + 1)$ -tuple of vertices contains at least one edge of  $\mathcal{H}$ , then  $\mathcal{E}_{k+1}^0 = \emptyset$ . In particular, taking  $\mathcal{H}$  as the complement of a hypergraph extremal for  $\text{ex}(n, \mathcal{K}_{k+1}^{(k)})$ , we obtain:

$$s'(n, k) \leq 1 - \frac{\text{ex}(n, \mathcal{K}_{k+1}^{(k)})}{\binom{n}{k}}.$$

As a consequence,

$$s'_k \leq 1 - \lim_{n \rightarrow \infty} \frac{\text{ex}(n, \mathcal{K}_{k+1}^{(k)})}{\binom{n}{k}}$$

where the limit exists for every fixed  $k$ , as proved in [5]. Hence, any lower bound on the Turán density of  $\mathcal{K}_{k+1}^{(k)}$  implies an upper bound on  $s'_k$ .

Note that an analogous implication in the opposite direction does not work: upper bounds on  $\text{ex}(n, \mathcal{K}_{k+1}^{(k)})$  do not imply lower bounds on  $s'(n, k)$ , or at least no evidence is found so far whether any function of the Turán numbers would yield a bound on  $s'(n, k)$  from below. On the other hand, applying the asymptotic result proved by Sidorenko [16], we obtain the following inequality.

**Corollary 13.** *As  $k$  gets large, we have  $s'_k \leq (1 + o(1)) \frac{\log k}{2k}$ .*

There are several further upper bounds which are more effective when  $k$  is small. For an overview of results of this kind and for their bibliographical data we refer to the survey [6]. De Caen’s conjecture [2], stating that  $O(1/k)$  is not a valid upper bound on  $1 - \lim_{n \rightarrow \infty} \frac{\text{ex}(n, \mathcal{K}_{k+1}^{(k)})}{\binom{n}{k}}$ , gives some indication on the possible limitations of this approach.

It cannot be guaranteed in general that the hypergraphs derived from the extremal ones for the Turán problem lead to constructions of  $(k - 1)$ -linear hypergraphs, hence they cannot automatically imply upper bounds on  $s_k$ . But this can be done if  $k = 2$ , and the following exact formula is valid.

**Theorem 14.** *For every  $n \geq 3$  we have  $s(n, 2) = s'(n, 2) = 1 - \left\lfloor \frac{n^2}{4} \right\rfloor / \binom{n}{2}$ , and therefore  $s_2 = s'_2 = 1/2$ . A hypergraph  $\mathcal{H} = (X, \mathcal{E})$  is extremal for  $s'(n, 2)$  if and only if  $\mathcal{E}_2$  is the complementary graph of the complete bipartite graph  $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ ; and for  $s(n, 2)$  the extremal hypergraph is unique up to isomorphism.*

**Proof.** For an upper bound, let  $\mathcal{H} = (X, \mathcal{E})$  consist of  $n$  vertices and two vertex-disjoint edges  $E_1, E_2$  with  $|E_1| = \lfloor n/2 \rfloor$  and  $|E_2| = \lceil n/2 \rceil$ . Then  $|\mathcal{E}_2| = \binom{n}{2} - \left\lfloor \frac{n^2}{4} \right\rfloor$  and  $\mathcal{E}_3^0 = \emptyset$ . Since  $\mathcal{H}$  is 1-linear, the upper bound follows for both  $s(n, 2)$  and  $s'(n, 2)$ . From the argument below, it will also turn out that this is the unique 1-linear hypergraph attaining equality.

To prove the lower bound, let  $\mathcal{H} = (X, \mathcal{E})$  be any hypergraph. Note that  $\mathcal{E}_2$  is just a graph; we denote its complement by  $G = (X, E)$ , i.e. an unordered vertex pair  $x_i x_j$  belongs to the edge set  $E$  of  $G$  if and only if  $\{x_i, x_j\} \notin \mathcal{E}_2$ . Then  $\mathcal{E}_3^0$  is the family of triangles ( $K_3$ -subgraphs) in  $G$ . As long as  $G$  is triangle-free, we have  $|\mathcal{E}_2| \geq \binom{n}{2} - \left\lfloor \frac{n^2}{4} \right\rfloor$  and the lower bound follows for  $\mathcal{H}$ , with equality if and only if  $G \simeq K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ . Assuming that  $|\mathcal{E}_2|$  is smaller, we have  $|E| > n^2/4$ .

We write the number  $|E|$  of edges in  $G$  in the form  $m = \frac{n^2}{4} + \ell$ ; hence  $\ell \geq 1$  is an integer if  $n$  is even, and  $\ell + \frac{1}{4} \geq 1$  is an integer if  $n$  is odd. It is well known that  $G$  has at least  $\frac{4m^2 - mn^2}{3n}$  triangles [12,13]. Thus,

$$|\mathcal{E}_3^0| \geq \frac{4m}{3n} \left( m - \frac{n^2}{4} \right) = \left( \frac{n}{3} + \frac{4\ell}{3n} \right) \ell > \frac{n\ell}{3}$$

and consequently

$$\begin{aligned} \frac{|\mathcal{E}_2|}{\binom{n}{2}} + \frac{|\mathcal{E}_3^0|}{\binom{n}{3}} &> \frac{\binom{n}{2} - \frac{n^2}{4} - \ell}{\binom{n}{2}} + \frac{\frac{n\ell}{3}}{\binom{n}{3}} \\ &= \frac{\binom{n}{2} - \frac{n^2}{4}}{\binom{n}{2}} + \frac{\frac{n\ell}{3} - \frac{(n-2)\ell}{3}}{\binom{n}{3}} \\ &\geq \frac{\binom{n}{2} - \left\lfloor \frac{n^2}{4} \right\rfloor}{\binom{n}{2}} + \frac{2\ell}{3\binom{n}{3}} - \frac{1}{4\binom{n}{2}}. \end{aligned}$$

This proves the stated inequality for all  $\ell \geq (n - 2)/8$ . Moreover, if  $n$  is even, the theorem follows for all  $\ell > 0$  because in that case we need not subtract  $1/4$  when moving from  $\frac{n^2}{4}$  to  $\lfloor \frac{n^2}{4} \rfloor$ .

For the rest of the proof, we assume that  $n$  is odd and  $0 < \ell < (n - 2)/8$ . Since  $\ell$  is relatively small,  $G$  must contain some vertex  $x$  of degree at most  $\frac{n-1}{2}$ , for otherwise the number of edges would be at least  $\frac{n(n+1)}{4}$ , yielding the contradiction  $\ell \geq \frac{n}{4}$ . Let now  $\ell' = \ell + \frac{1}{4}$ , that means  $m = \lfloor \frac{n^2}{4} \rfloor + \ell'$ , and consider the graph  $G' := G - x$ . This  $G'$  has  $n' = n - 1$  vertices and at least

$$m' := \frac{n^2}{4} + \ell - \frac{n - 1}{2} = \frac{(n - 1)^2 + 1}{4} + \ell = \frac{(n')^2}{4} + \ell'$$

edges. Therefore, by the theorem cited above,  $G'$  contains at least

$$\frac{4m'}{3n'} \left( m' - \frac{(n')^2}{4} \right) = \left( \frac{n - 1}{3} + \frac{4\ell'}{3n'} \right) \ell' > \frac{(n - 1)\ell'}{3}$$

triangles, which certainly is a lower bound on  $|\mathcal{E}_3^0|$ , too. Thus, with a slight modification of the computation above, we obtain that

$$\begin{aligned} \frac{|\mathcal{E}_2|}{\binom{n}{2}} + \frac{|\mathcal{E}_3^0|}{\binom{n}{3}} &> \frac{\binom{n}{2} - \lfloor \frac{n^2}{4} \rfloor - \ell'}{\binom{n}{2}} + \frac{\frac{(n-1)\ell'}{3}}{\binom{n}{3}} \\ &= \frac{\binom{n}{2} - \lfloor \frac{n^2}{4} \rfloor}{\binom{n}{2}} + \frac{\frac{(n-1)\ell'}{3} - \frac{(n-2)\ell'}{3}}{\binom{n}{3}} \\ &= \frac{\binom{n}{2} - \lfloor \frac{n^2}{4} \rfloor}{\binom{n}{2}} + \frac{\ell'}{3\binom{n}{3}}. \end{aligned}$$

This completes the proof of the theorem.

One of our referees remarks that the result can also be derived shortly from Razborov’s tight estimates [14] on the minimum number of triangles in a graph with a given number of vertices and edges. The proof of that result, however, can by no means be considered short, neither conceptually simple, as it applies the machinery of flag algebras.  $\square$

### 5. Concluding remarks

Motivated by a problem arising in chemistry/stoichiometry, we established asymptotically tight extremal results on geometric point sets and on finite set systems. Below we formulate some further problems and conjectures that remain open. *Geometry vs. hypergraph theory.* We proved corresponding asymptotic lower and upper bounds of the form  $\binom{n}{d+1} - \Theta(n^d)$  on the minimum number of affine simplices, for every set of  $n$  points in  $\mathbb{R}^d$  not containing any affine simplex of fewer than  $d + 1$  points. Our method was to put the problem in a more general context and to estimate an extremal function for a class of hypergraphs (called  $q$ -linear, implying the solution for geometric sets when  $q = d$ ). There remains a gap of order  $\Theta(n^d)$ , however, between the lower and upper bounds.

**Problem 15.** Given the integers  $n$  and  $d$ , determine the minimum number of affine simplices generated by  $n$  points in  $\mathbb{R}^d$ , no  $d$  of which lie on a  $(d - 2)$ -dimensional hyperplane.

**Problem 16.** Given the integers  $n$  and  $k$ , determine the minimum value of

$$|\mathcal{E}_k| + |\mathcal{E}_{k+1}^0|$$

taken over all  $(k - 1)$ -linear hypergraphs  $\mathcal{H} = (X, \mathcal{E})$  on  $n$  vertices.

**Problem 17.** For which values of  $k = d + 1$  is the minimum for hypergraphs in Problem 16 equal to that for point sets in  $\mathbb{R}^d$  in Problem 15, for all  $n > n_0(d)$ ?

For the case of  $d = 2$ , it was proved in [9] that the minimum number of affine simplices in  $\mathbb{R}^2$  determined by  $n \geq 8$  points is attained by placing the points on two lines: one of the lines contains  $n - 2$  of the points and the other line contains 3 points (and so their intersection point is also selected). That is, the minimum for Problem 15 with  $d = 2$  is

$$\binom{n - 2}{3} + \binom{n - 3}{2} + 1.$$

The construction implies the same upper bound for Problem 16 with  $k = 3$ , attained by the linear (that is, 2-linear) hypergraph with  $n$  vertices and two edges, one of size  $n - 2$  and the other of size 3. Moreover, the proof of the matching



lower bound in [9] gets through for linear hypergraphs as well, since it only applies modifications in the incidence structure, without any particular geometric assumptions. Thus, the minimum is the same for **Problem 15** with  $d = 2$  and **Problem 16** with  $k = 3$ .

It is not clear, however, whether the answer to **Problem 17** is positive or negative for  $d \geq 3$ . We note that the extremal construction of [17] cannot be applied for our problem to derive an upper bound on affine simplices in  $\mathbb{R}^3$ , because in [17] the points are arranged in two (equal or nearly equal) collinear sets. Nevertheless, the following conjecture looks easier than the exact determination of minimum.

**Conjecture 18.** *For every  $k \geq 3$  there exists a hypergraph  $\mathcal{H}$  of order  $n$  which is extremal for **Problem 16** and has  $O(n^{k-3})$  edges as  $n$  gets large.*

We note further that the upper bound on  $s'_k$  in **Corollary 13** tends to zero as  $k$  gets large, and at present we do not have any geometric constructions with the same property for  $s_k$ .

**Conjecture 19.** *There exists an integer  $k_0$  such that, for every  $k \geq k_0$ , we have  $s'_k < s_k$ .*

Perhaps the guess  $k_0 = 3$  is too brave, but we cannot disprove even that at present.

**Stoichiometry.** For the original problem originating from [7] in stoichiometry, our **Theorem 3** and **Corollary 4** imply:

- There are at least  $\binom{n}{d+1} - O(n^d) \approx \frac{n^{d+1}}{(d+1)!}$  minimal reactions among  $n$  species if the species are built up from  $d$  kinds of atoms (atomic particles), and if the number of species (molecules) forming any minimal reaction must be greater than  $d$ .
- For the first case previously unsolved, namely  $d = 3$ , the asymptotically tight lower bound is  $\binom{n}{4} - O(n^3) \approx \frac{n^4}{24}$ , if reactions with three or fewer species are not possible. (Especially parallel species, i.e. multiple doses are also excluded.)

## Acknowledgments

We thank the referees for their comments and proposed alternative approaches to some proofs. The research is supported in part by the Hungarian Scientific Research Fund, OTKA grant T-8 1493, and co-financed by the European Social Fund through the project TÁMOP-4.2.2.C-11/1/KONV-2012-0004 – National Research Center for Development and Market Introduction of Advanced Information and Communication Technologies.

## References

- [1] B. Bollobás, On generalized graphs, *Acta Math. Acad. Sci. Hungar.* 16 (1965) 447–452.
- [2] D. de Caen, The current status of Turán's problem on hypergraphs, in: P. Frankl, et al. (Eds.), *Extremal Problems for Finite Sets*, in: *Bolyai Society Mathematical Studies*, vol. 3, 1994, pp. 187–197.
- [3] Gy. Dósa, C. Laflamme, I. Szalkai, On the maximal and minimal number of bases and simple circuits in matroids and the extremal constructions, Preprint 046, Dept. Math. Univ. Veszprém, 1997.
- [4] Gy. Dósa, I. Szalkai, C. Laflamme, The maximum and minimum number of circuits and bases of matroids, *Pure Math. Appl. (PU.M.A.)* 15 (2004) 383–392.
- [5] Gy. Katona, T. Nemetz, M. Simonovits, Újabb bizonyítás a Turán-féle gráftételre és megjegyzések bizonyos általánosításaira, *Mat. Lapok* 15 (1964) 228–238. (in Hungarian).
- [6] P. Keevash, Hypergraph Turán problems, in: R. Chapman (Ed.), *Surveys in Combinatorics*, Cambridge University Press, 2011, pp. 83–140.
- [7] S. Kumar, Á. Pethő, Note on a combinatorial problem for the stoichiometry of chemical reactions, *Intern. Chem. Eng.* 25 (1985) 767–769.
- [8] C. Laflamme, I. Szalkai, Counting simplexes in  $\mathbb{R}^n$ , *Hung. J. Ind. Chem.* 23 (1995) 237–240.
- [9] C. Laflamme, I. Szalkai, Counting simplexes in  $\mathbb{R}^3$ , *Electron. J. Combin.* 5 (1) (1998) #R40. p. 11, Printed version in: *J. Combin.* 5 (1998), 597–607.
- [10] D. Lubell, A short proof of Sperner's lemma, *J. Combin. Theory* 1 (1966) 299.
- [11] L.D. Meshalkin, A generalization of Sperner's theorem on the number of subsets of a finite set, *Teor. Veroyatn. Primen.* 8 (1963) 219–220. (in Russian).
- [12] J.W. Moon, L. Moser, On a problem of Turán, *Magyar Tud. Akad. Mat. Kut. Int. Közl.* 7 (1962) 283–286.
- [13] E.A. Nordhaus, B.M. Steward, Triangles in an ordinary graph, *Canad. J. Math.* 15 (1963) 33–41.
- [14] A.A. Razborov, On the minimal density of triangles in graphs, *Combin. Probab. Comput.* 17 (2008) 603–618.
- [15] A. Sidorenko, What we know and what we do not know about Turán numbers, *Graphs Combin.* 11 (1995) 179–199.
- [16] A. Sidorenko, Upper bounds for Turán numbers, *J. Combin. Theory Ser. A* 77 (1997) 134–147.
- [17] B. Szalkai, I. Szalkai, Counting minimal reactions with specific conditions in  $\mathbb{R}^d$ , *J. Math. Chem.* 49 (2011) 1071–1085.
- [18] Zs. Tuza, Applications of the set-pair method in extremal hypergraph theory, in: P. Frankl, et al. (Eds.), *Extremal Problems for Finite Sets*, in: *Bolyai Society Mathematical Studies*, vol. 3, 1994, pp. 479–514.
- [19] Zs. Tuza, Applications of the set-pair method in extremal problems, II, in: D. Miklós, et al. (Eds.), *Combinatorics, Paul Erdős is Eighty*, in: *Bolyai Society Mathematical Studies*, vol. 2, 1996, pp. 459–490.
- [20] K. Yamamoto, Logarithmic order of free distributive lattices, *J. Math. Soc. Japan* 6 (1954) 343–354.
- [21] [https://en.wikipedia.org/wiki/Mikl%C3%B3s\\_Ybl](https://en.wikipedia.org/wiki/Mikl%C3%B3s_Ybl).